

AD-A093 726

NEW YORK UNIV N Y DEPT OF CHEMISTRY

F/6 7/4

PHOTOELECTRON EMISSION SPECTROSCOPY OF INORGANIC CATIONS IN AQUEOUS SOLUTIONS

DEC 80 P DELAHAY, K VON BURG, A DZIEDZIC

N00014-75-C-0397

UNCLASSIFIED


TR-13

NI

10/1
10/1



END
DATE
FILMED
2-84
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A093726

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0397

Task No. NR 051-258

TECHNICAL REPORT NO. 13 (New Series)

LEVEL II

Photoelectron Emission Spectroscopy of
Inorganic Cations in Aqueous Solution

by

Paul Delahay, Kathrin von Burg and Andrew Dziedzic

Prepared for Publication

in

Chemical Physics Letters

DTIC
S JAN 13 1981
E

New York University
Department of Chemistry
New York, NY

December 1980

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited

Accession For	
NTIS GS&I	X
DTIC TAB	
Unannounced	
Justification	
By	
Distribution	
Availability	
Dist	
A	

81 1 00 001

DOC FILE COPY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 13 (New Series -	2. GOVT ACCESSION NO. AD-A093	3. RECIPIENT'S CATALOG NUMBER 726
4. TITLE (and Subtitle) PHOTOELECTRON EMISSION SPECTROSCOPY OF INORGANIC CATIONS IN AQUEOUS SOLUTION.		5. TYPE OF REPORT & PERIOD COVERED (9) Technical Report
7. AUTHOR(s) Paul/Delahay, Kathrin/von Burg and Andrew/Dziedzic		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS New York University New York, NY 10003 (14) TR-13		8. CONTRACT OR GRANT NUMBER(s) (15) N00014-75-C-0397
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 051-258
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE December 1980
		13. NUMBER OF PAGES 22
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Submitted for publication in Chemical Physics Letters.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Aqueous solution Photoelectron spectroscopy Dielectric saturation Reorganization free energy Inorganic cation Standard potential Ionization potential Threshold energy Ligand field stabilization energy		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Threshold energies ($6.1 < E_t \leq 8.6$ eV) are determined for photoelectron emission by 16 inorganic cations in aqueous solution. E_t 's are correlated with gas-phase ionization potentials, solvation and reorganization free energies, standard reduction potentials and ligand field stabilization energies (five transition metals). Dielectric saturation is shown to drastically lower threshold energies.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-LP-014-6601Unclassified 406827
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

PHOTOELECTRON EMISSION SPECTROSCOPY OF INORGANIC CATIONS IN AQUEOUS SOLUTION

PAUL DELAHAY, KATHRIN von BURG and ANDREW DZIEDZIC

Department of Chemistry, New York University, 4 Washington Place, Room 514, New York, NY 10003, U.S.A.

Threshold energies ($6.1 < E_t \leq 8.6$ eV) are determined for photoelectron emission by 16 inorganic cations in aqueous solution. E_t 's are correlated with gas-phase ionization potentials, solvation and reorganization free energies, standard reduction potentials and ligand field stabilization energies (five transition metals). Dielectric saturation is shown to drastically lower threshold energies.

Threshold energies (ca. ± 0.1 eV) were recently obtained and interpreted [1,2] for photoelectron emission by various inorganic anions in aqueous solution. This work is extended to aqueous solutions of inorganic cations in the present paper.

1. Determination of threshold energies

Yields for photoelectron emission by solutions into water vapor were measured [1] as a function of photon energy (fig. 1). Anions with high threshold energies were selected to avoid emission by anions in the range of photon energies in which cations were investigated. Complexation [3] of the cation was avoided whenever possible. Hydrolysis [4] of the cation being photoionized and/or its photoionization product was minimized by addition of acid whenever necessary. Photoelectron emission by water [1] was totally negligible.

Threshold energies for emission were obtained [1,2] from emission spectra by extrapolation on the basis of the Brodsky-Tsarevsky theory [5]. A plot of the yield Y to the power 0.4 or 0.5 against photon energy is linear according to this theory, and extrapolation to zero yield gives the threshold energy

E_t . Selection of the proper exponent is discussed in [2] and [5]. The best plots were obtained by least square treatment of data, and the proper choice of exponent of Y was confirmed by calculation of correlation coefficients. The proportionality between the yield and the cation concentration was verified within ~ 10 percent for all cations (in general, 1, 0.5 and 0.2 M solutions) at constant anion concentration. This procedure insured that emission by the anion was negligible in the range of photon energies in which the yield was proportional to the cation concentration.


Threshold energies E_t are listed in Table 1. The slight variation of E_t from one anion to another for a given cation is essentially within experimental error. The standard deviation was 0.01 to 0.03 eV except for the 6 M acid solutions for which it was 0.06 to 0.08 eV because of a low signal (presumably because of scavenging of electrons by hydrogen ions). The effect of scavenging was also observed with reducible cations and is being investigated. E_t increased by 0.5 to 1 eV (vs. neutral solution) by addition of acid (≥ 2 M) for Mn^{2+} , Co^{2+} , Ni^{2+} . Addition of an alkali metal salt of the acid (≥ 2 M), instead of the acid, did not cause E_t to increase significantly. E_t for the acid was higher than for the alkali metal salt of the same anion as the acid, e.g., 9.08 eV for 6 M HCl vs. 8.77 eV for Cl^- in neutral solution (presumably because of removal of low-energy electrons by scavenging). This increase in E_t for Cl^- in 6 M HCl, for instance, allowed the determination of $E_t = 8.60$ eV for Co^{2+} without interference by the anion.

2. Threshold energy, ionization potential, reorganization and solvation free energies

The free energy for photoelectron emission by an aqueous solution of cation $M^{2+}(aq)$ will be correlated with the free energy for gas-phase ionization of $M^{2+}(g)$. The reference level of free energies must be changed

for that purpose. The problem is discussed in detail by Noyes [7,8], and only essential points will be mentioned. The surface potential at the solution-water vapor interface (~ 0.1 to 0.2 V for water [9]) is neglected here.

Free energies must be changed from their conventional values (i.e., in tables [10]) to the reference level of an electron in the gas phase at infinity at the same outer potential [11] as the solution. Consider the reactions,



for which (g) and (aq) specify the gas phase and solution, respectively. The free energies are (fig. 2): the conventional value ΔG for (1); $\Delta G_H = 4.50$ eV [8] for (2); $\Delta G + \Delta G_H$ for (3). Likewise, the conventional free energies of solvation of $M^{Z+}(aq)$ and $M^{(Z+1)+}(aq)$ ΔG_C^Z and ΔG_C^{Z+1} are converted to $\Delta G_S^Z = \Delta G_C^Z + Z \Delta G_H$ and $\Delta G_S^{Z+1} = \Delta G_C^{Z+1} + (Z+1) \Delta G_H$, respectively.

Photoionization of $M^{Z+}(aq)$ yields initially the species $M^{(Z+1)+}(aq^*)$ under conditions governed by the Franck-Condon principle. The solvent about this ion undergoes reorganization to yield the ion $M^{(Z+1)+}(aq)$ in a process characterized by the reorganization free energy ΔG_r^{Z+1} .

Two basic equations follow from fig. 2:

$$\Delta G_i^Z(g) = \Delta G_S^Z - \Delta G_S^{Z+1} + \Delta G + \Delta G_H \quad (4)$$

$$\Delta G_e^Z(aq) = \Delta G_i^Z(g) + \Delta G_S^{Z+1} - \Delta G_S^Z - \Delta G_r^{Z+1} \quad (5)$$

where $\Delta G_e^Z(aq)$ and $\Delta G_i^Z(g)$ are, respectively, the free energies for photoelectron emission by the solution and gas-phase ionization. A third equation follows from fig. 2 or can be obtained by combination of (4) and (5),

$$\Delta G_e^Z(\text{aq}) = \Delta G + \Delta G_H - \Delta G_r^{Z+1}. \quad (6)$$

One has $\Delta G_e^Z(\text{aq}) = E_t$ to a good approximation since measurements yield free energies for photoelectron emission. $\Delta G_i^Z(\text{g})$ can be calculated from (4), e.g., 29.81 and 30.51 eV, respectively, for Cr^{2+} and Fe^{2+} (data from Table 1). The necessary ΔG_s^{Z+1} , however, is generally not available and therefore we shall use ionization potentials I^Z instead of $\Delta G_i^Z(\text{g})$ in sec. 3. The I^Z 's are enthalpies, but we set $\Delta G_i^Z(\text{g}) = I^Z$ and thus neglect a minor entropy correction.

Reorganization free energies ΔG_r^{Z+1} can be computed from (5), e.g., -2.05 and -2.11 eV, respectively, for Cr^{2+} and Fe^{2+} on the basis of the $\Delta G_i^2(\text{g})$ calculated above from (4) for these two cations. Since $\Delta G_s^3 = 45.04$ and 44.87 eV for Cr^{3+} and Fe^{3+} , respectively, one has $|G_r^3| \ll |\Delta G_s^3|$ for Cr^{2+} and Fe^{2+} . Hence, the ion $M^{(Z+1)+}(\text{aq}^*)$ produced by vertical transition in the photoionization of $M^{Z+}(\text{aq})$ is energetically not very different from $M^{(Z+1)+}(\text{aq})$.

The orientation polarization of the solvent about $M^{(Z+1)+}(\text{aq}^*)$ is the same as for $M^{Z+}(\text{aq})$ (Franck-Condon principle). Conversely, the electronic polarization of the solvent about $M^{(Z+1)+}(\text{aq}^*)$ is the same as for $M^{(Z+1)+}(\text{aq})$ except for a secondary effect arising from the different orientation polarizations for these two ions. Since the free energies of $M^{(Z+1)+}(\text{aq}^*)$ and $M^{(Z+1)+}(\text{aq})$ are quite close, as shown above, the orientation polarization about $M^{Z+}(\text{aq})$ and $M^{(Z+1)+}(\text{aq})$ must not be very different. This can be the case only if there is strong dielectric saturation of the solvent about these ions.

The preceding conclusion is in complete agreement with Noyes [7,8] who calculated effective dielectric constants from solvation free energies and crystallographic radii by application of the Born equation. The resulting

effective dielectric constants (well below 10) indicated strong dielectric saturation. It should be noted that the foregoing analysis, based on eq. (5) does not involve the Born model of solvation.

3. Threshold energy and dielectric saturation

The calculation of ΔG_r^{z+1} in sec. 2 was restricted to Cr^{3+} and Fe^{3+} because of the lack of complete data for other cations. A method will be developed in this section by which the essential role of dielectric saturation will be established for all the cations of Table 1. The method involves the Born model of solvation, but the approximation inherent to the use of this model does not affect the conclusion to be drawn. Thus,

$$\Delta G_r^{z+1} = \Delta G_s^{z+1} + p_o^z + p_e^{z+1}(p_o^z), \quad (7)$$

where $p_o^z (> 0)$ is the orientation polarization about $M^{(z+1)+}(\text{aq}^*)$ and $p_e^{z+1}(p_o^z)$ the corresponding electronic polarization (see above). A very small correction (< 0.05 eV) is neglected in (7) for the change of solvation free energy of an electrically neutral species [7] as the cavity radius changes from the value for p_o^z to that for p_o^{z+1} .

One has,

$$|\Delta G_s^z| < p_o^z + p_e^{z+1}(p_o^z) < |\Delta G_s^{z+1}|, \quad (8)$$

and consequently we set

$$p_o^z + p_e^{z+1}(p_o^z) = -\beta \Delta G_s^z \quad (9)$$

where the coefficient $\beta (> 1)$ is to be determined from experimental data and calculated from a model. Equation (5), (7) and (9) yield,

$$\Delta G_e^z(\text{aq}) = \Delta G_i^z(\text{g}) + (\beta - 1) \Delta G_s^z. \quad (10)$$

The coefficient β can be calculated from the Born equation. Thus,

$$\beta = [(z+1)/z]^2 (1 - \epsilon_0^{-1}) / (1 - \epsilon^{-1})^{-1} + (\epsilon_0^{-1} - \epsilon^{-1}) / (1 - \epsilon^{-1}), \quad (11)$$

where ϵ_0 is the optical dielectric constant of water and ϵ its static dielectric constant in the absence of dielectric saturation. One computes

from (11), $\beta = 2.33, 1.55, 1.34$ for $z = 1, 2, 3$, respectively, in the absence of dielectric saturation ($\epsilon = 78.36$). Conversely, one has $\beta = [(z+1)/z]^2$ for complete saturation ($\epsilon = \infty$), that is, $\beta = 4, 2.25, 1.78$, respectively, for $z = 1, 2, 3$.

Values of β calculated from (10) with $\Delta G_i^z(g) = I^z$ (see above) are listed in Table 1. The ionization potentials I^z (from [4] except for Cr^{2+} and Fe^{3+} [6]) are more recent and in some cases slightly different from those used by Noyes (from [10]) in the calculation of conventional solvation free energies. This minor inconsistency is of no consequence in the interpretation of β . The β -values of Table 1 are close to those calculated from (11) for complete dielectric saturation. Equation (11) is approximate (Born model), and partial hydrolysis and/or complexation may cause β to be higher than expected for the ionic charge $z+$. Moreover, the value of β is not very sensitive to $\Delta G_e^z(\text{aq})$. Despite these reservations, one can safely conclude that there is strong dielectric saturation of water about the cations of Table 1. Reorganization free energies would be much higher in the absence of dielectric saturation, and the threshold energies of cations in aqueous solution are drastically lowered by strong dielectric saturation of the solvent about the cations.

4. Threshold energy, standard potential and reorganization free energy

The free energy ΔG in (6) pertains to reaction (1) which is written as an oxidation. Hence, $\Delta G = |e|E^0$ (in eV) where E^0 is the standard reduction potential for the $\text{M}^{z+1}(\text{aq})/\text{M}^{z+}(\text{aq})$ couple. Equation (6) becomes (in eV)

$$\Delta G_e^z(\text{aq}) = |e|E^0 + 4.50 - \Delta G_r^{z+1}, \quad (12)$$

where $\Delta G_H = 4.50$ eV was introduced (sec. 2). The free energy (ca. 0.1 to 0.2 eV [9]) for the surface potential should be added on the right hand side of (12).

Equation (12) clearly shows the relationship between the free energies $\Delta G_e^Z(\text{aq})$ and ΔG for photoelectron emission and thermodynamics in the oxidation of $M^{Z+}(\text{aq})$ to $M^{Z+1}(\text{aq})$. ΔG_H arises from the change of reference levels for free energies, and $-\Delta G_r^{Z+1}$ is the additional free energy required by the Franck-Condon principle. The negative of the reorganization free energy can be regarded as the counterpart of the overvoltage in the electrochemical oxidation of $M^{Z+}(\text{aq})$.

Reorganization free energies calculated from (12) are nearly the same (-2.05 to -2.25 eV) for the five cations of fig. 3 (E^0 's from [4], [12]). Since ΔG_r^3 is nearly constant, a plot of E_t against $|e| E^0$ [2,13] is linear, to a first approximation, and has a unit slope (fig. 3). Standard reduction potentials can be deduced from such a plot and E_t (for chemically similar cations) for species that are unstable in aqueous solution, e.g., $E^0 \approx 1.75 \text{ V}$ for $\text{Ni}^{3+}/\text{Ni}^{2+}$ in 6 M HCl (using $\Delta G_r = -2.1 \text{ eV}$).

5. Threshold energy, ionization potential and ligand field stabilization energies (LFSE)

The $M^{2+}(\text{aq})$ and $M^{3+}(\text{aq})$ ions of V, Cr, Mn, Fe, Co exist as hexaquo ions [4]. $\Delta G_e^Z(\text{aq})$ therefore include the difference ΔE_f between the LFSE's for $M^{3+}(\text{aq})$ and $M^{2+}(\text{aq})$ ions in an octahedral field. I^2 for these metals obviously does not include ΔE_f . Hence the correlation between I^2 and E_t is partially masked (for Cr, in particular, fig. 4) by the LFSE effect. The correlation, however, is evident when I^2 and $E_t + \Delta E_f$ are compared. Thus, $I^2 - (E_t + \Delta E_f)$ and $\Delta G_s^2 - \Delta G_s^3 + \Delta G_r^3 - \Delta E_f$ (cf. eq. (5)) are nearly constant from V to Co whereas $I^2 - E_t$ is not.

Acknowledgment

This work was supported by the Office of Naval Research and the National Science Foundation.

References

- [1]I. Watanabe, J. B. Flanagan and P. Delahay, J. Chem. Phys. 73 (1980) 2057.
- [2]K. von Burg and P. Delahay, Chem. Phys. Lett., submitted.
- [3]L. G. Sillen and A. E. Martell, Stability Constants, Special Publication No. 17, Supplement No. 1 (The Chemical Society, London, 1971).
- [4]F. A. Cotton and G. Wilkinson, Advanced Inorganic Chemistry, 4th ed. (Interscience-Wiley, New York, 1980).
- [5]A. M. Brodsky and A. V. Tsarevsky, J. Chem. Soc. Faraday Trans. II 72 (1976) 1781.
- [6]Handbook of Chemistry and Physics (CRC Press, Cleveland, Ohio, 1975, p. E-68).
- [7]R. M. Noyes, J. Am. Chem. Soc. 84 (1962) 513.
- [8]R. M. Noyes, J. Am. Chem. Soc. 86 (1964) 971.
- [9]J. E. B. Randles in Advances in Electrochemistry and Electrochemical Engineering, P. Delahay and C. W. Tobias, editors, vol. 3 (Wiley-Interscience, New York, 1963, pp. 1-30).
- [10]Selected Values of Chemical Thermodynamic Properties, Circular No. 500 of the National Bureau of Standards (U. S. Government Printing Office, Washington, D. C., 1952).
- [11]J. Koryta, J. Dvorak and V. Bohackova, Electrochemistry (Methuen, London, 1970, pp. 140-149).
- [12]W. M. Latimer, Oxidation Potentials, 2nd ed. (Prentice-Hall, New York, 1952).
- [13]R. E. Ballard, Chem. Phys. Lett. 42 (1976) 97.
- [14]T. M. Dunn, D. S. McClure, R. G. Pearson, Crystal Field Theory (Harper and Row, New York, 1965, p. 82).

Table 1

Threshold energies, coefficient β and relevant data

	E_t eV	I^Z [4,6] eV	$-\Delta G_s^Z$ eV	β
Ag^+	7.60 (ClO_4^-), 7.52 (ClO_4^- , 5 M $HClO_4$)	21.48	4.95	3.80
Tl^+	7.40 (F^-), 7.46 (ClO_4^- , 5 M $HClO_4$)	20.42	3.55	4.67
V^{2+}	6.38 (SO_4^{2-} , 1 M H_2SO_4)	29.31	19.14	2.20
Cr^{2+}	6.14 (Cl^- , 1 M HCl)	30.95	19.32	2.28
Mn^{2+}	8.08 (Cl^- , 2 M HCl)	33.70	18.96	2.35
Fe^{2+}	7.35 (Cl^- , 1 M HCl), 7.38 (SO_4^{2-})	30.64	19.63	2.18
Co^{2+}	8.60 (Cl^- , 6 M HCl)	33.49	20.88	2.19
Ni^{2+}	8.35 (Cl^- , 6 M HCl)	36.16	21.40	2.30
Cu^{2+}	7.83 (ClO_4^- , 1 M $HClO_4$), 7.84 (SO_4^{2-})	36.74	21.55	2.34
Sn^{2+}	7.21 (Cl^-), 7.42 (ClO_4^- , 6 M $HClO_4$)	30.49	16.14	2.44
Pb^{2+}	7.23 (ClO_4^-), 7.10 (ClO_4^- , 6 M $HClO_4$)	31.92	15.49	2.59
In^{3+}	7.15 (ClO_4^-)	54.42	42.57	2.11
Tl^{3+}	6.90 (Cl^-)	43.25	42.01	1.87
V^{3+}	7.06 (Cl^-)	47.7	44.2	1.92
Cr^{3+}	7.33 (Cl^-), 7.27 (SO_4^{2-})	50.8	45.04	1.97
Fe^{3+}	7.03 (Cl^-), 7.05 (Cl^- , 6 M $HClO_4$)	56.8	44.87	2.11

See sec. 1 for standard deviation. ΔG_2^3 for V^{3+} calculated from (4) with $\Delta G = -0.25$ eV (sec. 4).

Captions to Figures

Fig. 1. Photoelectron emission spectra of Fe^{2+} . (A) 1 M FeSO_4 ; (B) 0.5 M FeSO_4 + 0.5 M Li_2SO_4 ; (C) 0.2 M FeSO_4 + 0.8 M Li_2SO_4 . Photoelectron emission only by Fe^{2+} below the threshold energy ($E_t = 8.65$ eV) for SO_4^{2-} .

Fig. 2 Free energy diagram. Upward and downward arrows, respectively, for positive and negative free energies. The superscripts z and $z+1$ denote the positive ionic charge. The symbols (g) and (aq) refer to gas phase and solution, respectively. Free energies: $\Delta G^z(\text{g})$, $\Delta G^{z+1}(\text{g})$, $\Delta G^z(\text{aq})$, $\Delta G^{z+1}(\text{aq})$ of ions; ΔG for reaction (1); ΔG_H for (2); ΔG_C^z and ΔG_C^{z+1} conventional values for solvation; ΔG_S^z , ΔG_S^{z+1} values for solvation corrected for ΔG_H ; ΔG_r^{z+1} reorganization; $\Delta G_i^z(\text{g})$ gas-phase ionization; $\Delta G_e^z(\text{aq})$ photoelectron emission by solution.

Fig. 3 Plot of threshold energy E_t and reorganization free energy ΔG_r^3 against $|e|E^0$. Standard reduction potentials E^0 from [4,12]. The E^0 's for $\text{Mn}^{3+}/\text{Mn}^{2+}$ and $\text{Co}^{3+}/\text{Co}^{2+}$ are approximate (ca. ± 0.05 V). The symbol M_*^{3+} stands for $\text{M}^{3+}(\text{aq}^*)$.

Fig. 4 Correlation between the ionization potential I^2 and the threshold energy E_t corrected for the difference ΔE_f between the LFSE's for M^{3+} and M^{2+} in an octahedral field. Values of ΔE_f from [14].




FIG. 1




FIG. 2




FIG. 3




FIG. 4

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Area Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Western Regional Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Paul Delahay Department of Chemistry New York University New York, New York 10003	1	Dr. P. J. Hendra Department of Chemistry University of Southampton Southampton SO9 5NH United Kingdom	1
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 44106	1	Dr. Sam Perone Department of Chemistry Purdue University West Lafayette, Indiana 47907	1
Dr. D. N. Bennion Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
Dr. J. J. Auborn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research & Technology Department Naval Surface Weapons Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. T. Katan Lockheed Missiles & Space Co, Inc. P.O. Box 504 Sunnyvale, California 94088	1	Dr. J. Boechler Electrochimica Corporation Attention: Technical Library 2485 Charleston Road Mountain View, California 94040	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063	1
Dr. B. Brummer EIC Incorporated 55 Chapel Street Newton, Massachusetts 02158	1	Dr. F. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1	Dr. R. P. Van Duyne Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. B. Stanley Pons Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6G 2G2	1
Larry E. Flew Naval Weapons Support Center Code 30736, Building 2906 Crane, Indiana 47522	1	Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824	1
S. Ruby DOE (STOR) 600 E Street Washington, D.C. 20545	1	Dr. R. David Rauh EIC Corporation 55 Chapel Street Newton, Massachusetts 02158	1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	1	Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1	Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton SO9 5NH England	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1	Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California	1	Dr. R. A. Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910	1	Mr. James R. Moden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. R. Novak Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Dr. John Kincaid Department of the Navy Strategic Systems Project Office Room 901 Washington, DC 20376	1
Dr. John F. Houlihan Shenango Valley Campus Pennsylvania State University Sharon, Pennsylvania 16146	1	M. L. Robertson Manager, Electrochemical Power Sonics Division Naval Weapons Support Center Crane, Indiana 47522	1
Dr. M. G. Sceats Department of Chemistry University of Rochester Rochester, New York 14627	1	Dr. Elton Cairns Energy & Environment Division Lawrence Berkeley Laboratory University of California Berkeley, California 94720	1
Dr. D. F. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201	1	Dr. Bernard Spielvogel U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709	1
Dr. D. H. Whitmore Department of Materials Science Northwestern University Evanston, Illinois 60201	1	Dr. Denton Elliott Air Force Office of Scientific Research Bldg. 104 Bolling AFB Washington, DC 20332	1
Dr. Alan Bewick Department of Chemistry The University Southampton, SO9 5NH England	1		
Dr. A. Hiny NAVSEA-5433 NC #4 2541 Jefferson Davis Highway Arlington, Virginia 20362	1		

TECHNICAL REPORT DISTRIBUTION LIST, 051B

	<u>No. Copies</u>		<u>No. Copies</u>
Professor K. Wilson Department of Chemistry, B-014 University of California, San Diego La Jolla, California 92093	1	Dr. B. Vonnegut State University of New York Earth Sciences Building 1400 Washington Avenue Albany, New York 12203	1
Professor C. A. Angell Department of Chemistry Purdue University West Lafayette, Indiana 47907	1	Dr. Hank Loos Laguna Research Laboratory 21421 Stans Lane Laguna Beach, California 92651	1
Professor P. Meijer Department of Physics Catholic University of America Washington, D.C. 20064	1	Dr. John Latham University of Manchester Institute of Science & Technology P.O. Box 88 Manchester, England M601QD	1
Dr. S. Greer Chemistry Department University of Maryland College Park, Maryland 20742	1		
Professor P. Delahay New York University 100 Washington Square East New York, New York 10003	1		
Dr. T. Ashworth Department of Physics South Dakota School of Mines & Technology Rapid City, South Dakota 57701	1		
Dr. G. Gross New Mexico Institute of Mining & Technology Socorro, New Mexico 87801	1		
Dr. J. Kassner Space Science Research Center University of Missouri - Rolla Rolla, Missouri 65401	1		
Dr. J. Telford University of Nevada System Desert Research Institute Lab of Atmospheric Physics Reno, Nevada 89507	1		

